

## **Optimisation of Vitamin D Status for Enhanced Immuno-protection against Covid-19**

**Dr. Daniel M. McCartney, Lecturer in Human Nutrition & Dietetics, School of Biological and Health Sciences, College of Sciences and Health, Technological University of Dublin City Campus.**

**Dr. Declan G. Byrne, Consultant General Physician, Department of Internal Medicine, St. James's Hospital, Dublin 8 and Clinical Senior Lecturer, Dept. of Clinical Medicine, School of Medicine, Trinity College Dublin.**

### **Healthcare Professionals Summary Fact Sheet**

#### ***Background***

Vitamin D can be made in our skin, taken in the diet or derived from supplements. Historically, humans received most of their vitamin D from sunshine exposure during the Summer months, with diet contributing only very modest amounts in relation to our overall needs. As our sunlight exposure has declined over recent decades, we have become increasingly reliant on inadequate dietary sources to meet our physiological demands for vitamin D<sup>1</sup>.

Consequently, most people cannot now meet their vitamin D requirements from diet alone without taking supplements<sup>2,3</sup>.

#### ***Prevalence of Vitamin D Deficiency***

It has been shown that many people in Ireland, particularly in late Winter and early Spring, have low serum levels of 25-hydroxy vitamin D (25(OH)D)<sup>4</sup>, the metabolite used to assess vitamin D status. While many experts contend that optimal serum levels are those above 75-80nmol/l<sup>5</sup>, there is general agreement that levels below 50nmol/l constitute deficiency, and that levels less than 30nmol/l signify severe deficiency<sup>6</sup>.

Older adults are much more likely to have low serum 25(OH) D than younger adults. In Ireland, the National Adults Nutrition Survey (NANS) showed that 36% of adults aged 50-64 years, and 44% of adults aged 65-84 years had serum vitamin D levels less than 50nmol/l on a year-round basis, while the prevalence of deficiency in those age groups rose to 55% and 48% respectively in winter<sup>4</sup>. It has further been shown that hospital inpatients and nursing home residents in Ireland have serum vitamin D levels which are even lower than those of community-dwelling older adults.<sup>7</sup>

Other groups who are likely to have low serum 25(OH)D levels include:

- those with darker skin
- those who avoid sun exposure during Summer
- vegetarians and vegans
- those who are overweight or obese
- smokers

### ***Vitamin D and Respiratory Infection***

Epidemiological studies, including several meta-analyses, have shown that people with low vitamin D levels have a higher risk of acute respiratory tract infection and community-acquired pneumonia<sup>8,9</sup>.

While these data do not necessarily infer causality, multiple molecular mechanisms have been identified by which vitamin D deficiency impairs resistance to viral respiratory tract infection<sup>10,11</sup>.

There are also a significant number of studies, including several meta-analyses, which have indicated that vitamin D supplementation may reduce the likelihood of acute respiratory tract infection, and decrease its severity and duration where such infection does occur<sup>12,13</sup>.

These respiratory tract infections may include Covid-19.

### ***Proposed Protective Mechanisms against Covid-19***

In this regard, vitamin D supplementation has been shown to suppress CD26<sup>14</sup>, a cell surface receptor which is thought to facilitate entry of the Covid-19 virus into the host cell<sup>15</sup>.

There is also good evidence that enhanced vitamin D status may protect against the critical immunological sequelae which are thought to elicit poorer clinical outcome in Covid-19 infection. These include prolonged interferon-gamma response<sup>11</sup>, and persistent interleukin 6 elevation, a negative prognostic indicator in acutely-ill pneumonia patients<sup>16</sup>, including those with Covid-19.

### ***Supplementation Guidelines***

The Food Safety Authority of Ireland (FSAI), the European Food Safety Authority (EFSA) and many other international agencies already recommend that both adults and children supplement with vitamin D. While the FSAI recommends a supplement of 10 micrograms per day for older adults<sup>3</sup>, most countries in Europe now recommend 15-20 micrograms per day for these older age groups<sup>6</sup>. In the US, the Institute of Medicine (IoM) has suggested an oral intake of 20 micrograms per day for those aged 70 years and over<sup>17</sup>, while the Endocrine Society has proposed an intake of 37.5-50 micrograms per day for older adults to optimise their serum 25(OH)D levels<sup>5</sup>.

Research has indicated that Irish adults require ~25-30 micrograms of vitamin D per day to reliably maintain their serum levels above 50nmol/l on a year round basis<sup>18,19</sup>, while numerous studies have also demonstrated the safety of supplementing with vitamin D at doses of 20-50 micrograms per day (800-2000 iu per day)<sup>20-22</sup>. The latter finding is perhaps unsurprising as this is an oral dose which is lower than the amount of vitamin D typically synthesised by skin exposure to sunshine on a Summer's day<sup>23,24</sup>.

## **Conclusions and Recommendations**

For those with serum levels less than 50nmol/l, each additional microgram of intake per day from supplements elicits a serum 25(OH)D increase of about 1nmol/l<sup>25</sup>. This implies that supplementing at 20-50 micrograms per day will raise serum 25(OH)D to the levels above 50nmol/l which have been associated with reduced risk of viral respiratory tract infection.

It is proposed that such vitamin D supplementation may enhance resistance to Covid-19 infection, or limit its damaging immunological sequelae and improve clinical prognosis in those who do become infected.

Consequently, all older adults, hospital inpatients and nursing home residents should be supplemented with a *minimum* 20 microgram daily dose of vitamin D. This guidance should also be prioritised in other vulnerable groups including those with diabetes mellitus or compromised immune function, individuals with limited habitual sun exposure, those with darker skin, vegans, vegetarians, smokers and those who are overweight or obese. It is further recommended that given their potential occupational exposure, this advice be urgently promoted to all healthcare professionals, and ultimately extended to the remainder of the adult population to mitigate their risk of Covid-19 infection.

## **References**

1. Rajakumar K, Greenspan SL, Thomas SB, Holick MF. SOLAR ultraviolet radiation and vitamin D: a historical perspective. *Am J Public Health*. 2007; 97:1746-54.
2. Veugelers PJ, Pham TM, Ekwaru JP. Optimal Vitamin D Supplementation Doses that Minimize the Risk for Both Low and High Serum 25-Hydroxyvitamin D Concentrations in the General Population. *Nutrients*. 2015; 7:10189-208. doi: 10.3390/nu7125527.
3. Food Safety Authority of Ireland. *Scientific Recommendations for Healthy Eating Guidelines in Ireland*. Dublin: Food Safety Authority of Ireland, 2011.
4. Cashman KD, Muldowney S, McNulty B, Nugent A, FitzGerald AP, Kiely M, Walton J, Gibney MJ, Flynn A. Vitamin D status of Irish adults: findings from the National Adult Nutrition Survey. *Br J Nutr*. 2013; 109:1248-56. doi: 10.1017/S0007114512003212.
5. Holick MF, Binkley NC, Bischoff-Ferrari HA, Gordon CM, Hanley DA, Heaney RP, Murad MH, Weaver CM & Endocrine Society. Evaluation, treatment, and prevention of vitamin D deficiency: an Endocrine Society clinical practice guideline. *Journal of Clinical Endocrinology and Metabolism*. 2011; 96:1911–1930.
6. Lips P, Cashman KD, Lamberg-Allardt C, Bischoff-Ferrari HA, Obermayer-Pietsch B, Bianchi ML, Stepan J, El-Hajj Fuleihan G, Bouillon R. Current vitamin D status in European and Middle East countries and strategies to prevent vitamin D deficiency: a position statement of the European Calcified Tissue Society. *Eur J Endocrinol*. 2019; 180:P23-P54. doi: 10.1530/EJE-18-0736.
7. Griffin TP, Wall D, Blake L, Griffin DG, Robinson SM, Bell M, Mulkerrin EC, O'Shea PM. Vitamin D status of adults in the community, in outpatient clinics, in hospital and in nursing homes in the West of Ireland. *J Gerontol A Biol Sci Med Sci*. 2020 Jan 14. pii: glaa010. doi: 10.1093/gerona/glaa010.

8. Martineau AR, Jolliffe DA, Hooper RL, Greenberg L, Aloia JF, Bergman P, Dubnov-Raz G, Esposito S, Ganmaa D, Ginde AA, Goodall EC, Grant CC, Griffiths CJ, Janssens W, Laaksi I, Manaseki-Holland S, Mauger D, Murdoch DR, Neale R, Rees JR, Simpson S Jr, Stelmach I, Kumar GT, Urashima M, Camargo CA Jr. Vitamin D supplementation to prevent acute respiratory tract infections: systematic review and meta-analysis of individual participant data. *BMJ*. 2017; 356:i6583. doi: 10.1136/bmj.i6583.
9. Zhou YF, Luo BA, Qin LL. The association between vitamin D deficiency and community-acquired pneumonia: A meta-analysis of observational studies. *Medicine (Baltimore)*. 2019; 98:e17252. doi: 10.1097/MD.00000000000017252.
10. Telcian AG, Zdrengea MT, Edwards MR, Laza-Stanca V, Mallia P, Johnston SL, Stanciu LA. Vitamin D increases the antiviral activity of bronchial epithelial cells in vitro. *Antiviral Res*. 2017; 137:93-101. doi: 10.1016/j.antiviral.2016.11.004.
11. Zdrengea MT, Makrinioti H, Bagacean C, Bush A, Johnston SL, Stanciu LA. Vitamin D modulation of innate immune responses to respiratory viral infections. *Rev Med Virol*. 2017; 27(1). doi: 10.1002/rmv.1909.
12. Rejnmark L, Bislev LS, Cashman KD, Eiríksdóttir G, Gaksch M, Gröbler M, Grimnes G, Gudnason V, Lips P, Pilz S, van Schoor NM, Kiely M, Jorde R. Non-skeletal health effects of vitamin D supplementation: A systematic review on findings from meta-analyses summarizing trial data. *PLoS One*. 2017; 12:e0180512. doi: 10.1371/journal.pone.0180512. eCollection 2017.
13. Autier P, Mullie P, Macacu A, Dragomir M, Boniol M, Coppens K, Pizot C, Boniol M. Effect of vitamin D supplementation on non-skeletal disorders: a systematic review of meta-analyses and randomised trials. *Lancet Diabetes Endocrinol*. 2017; 5:986-1004. doi: 10.1016/S2213-8587(17)30357-1.
14. Skariyachan S, Challapilli SB, Packirisamy S, Kumargowda ST, Sridhar VS. Recent Aspects on the Pathogenesis Mechanism, Animal Models and Novel Therapeutic Interventions for Middle East Respiratory Syndrome Coronavirus Infections. *Front Microbiol*. 2019; 10:569. doi: 10.3389/fmicb.2019.00569.
15. Vankadari N, Wilce JA. Emerging WuHan (COVID-19) coronavirus: glycan shield and structure prediction of spike glycoprotein and its interaction with human CD26. *Emerg Microbes Infect*. 2020; 9:601-604. doi: 10.1080/22221751.2020.1739565.
16. Miroliaee AE, Salamzadeh J, Shokouhi S, Sahraei Z. The study of vitamin D administration effect on CRP and Interleukin-6 as prognostic biomarkers of ventilator associated pneumonia. *J Crit Care*. 2018; 44:300-305. doi: 10.1016/j.jcrc.2017.08.040.
17. Institute of Medicine. *Dietary Reference Intakes for Calcium and Vitamin D*. Washington DC: The National Academies Press, 2011.
18. Cashman KD, Hill TR, Lucey AJ, Taylor N, Seamans KM, Muldowney S, Fitzgerald AP, Flynn A, Barnes MS, Horigan G, Bonham MP, Duffy EM, Strain JJ, Wallace JM, Kiely M. Estimation of the dietary requirement for vitamin D in healthy adults. *Am J Clin Nutr*. 2008; 88:1535-42. doi: 10.3945/ajcn.2008.26594.
19. Cashman KD, Wallace JM, Horigan G, Hill TR, Barnes MS, Lucey AJ, Bonham MP, Taylor N, Duffy EM, Seamans K, Muldowney S, Fitzgerald AP, Flynn A, Strain JJ, Kiely M. Estimation of the dietary requirement for vitamin D in free-living adults  $\geq 64$  y of age. *Am J Clin Nutr*. 2009; 89:1366-74. doi: 10.3945/ajcn.2008.27334.

20. Vieth R, Kimball S, Hu A, Walfish PG. Randomized comparison of the effects of the vitamin D3 adequate intake versus 100 mcg (4000 IU) per day on biochemical responses and the wellbeing of patients. *Nutr J.* 2004; 3:8.
21. Bischoff-Ferrari HA, Shao A, Dawson-Hughes B, Hathcock J, Giovannucci E, Willett WC. Benefit-risk assessment of vitamin D supplementation. *Osteoporos Int.* 2010; 21:1121-32. doi: 10.1007/s00198-009-1119-3.
22. Hathcock JN, Shao A, Vieth R, Heaney R. Risk assessment for vitamin D. *Am J Clin Nutr.* 2007; 85:6-18.
23. Barger-Lux MJ, Heaney RP. Effects of above average summer sun exposure on serum 25-hydroxyvitamin D and calcium absorption. *J Clin Endocrinol Metab.* 2002; 87:4952-6.
24. Heaney RP, Armas LA. Quantifying the vitamin D economy. *Nutr Rev.* 2015; 73:51-67. doi: 10.1093/nutrit/nuu004.
25. Vieth R. Why the optimal requirement for Vitamin D3 is probably much higher than what is officially recommended for adults. *J Steroid Biochem Mol Biol.* 2004; 89-90:575-9.